

## Factors Associated with Measles Occurrence among Children Aged 6 Months to 10 Years in Ludimbi Health Zone, Lukula, Democratic Republic of the Congo, 2016–2022: A Case-Control Study.

Théophile Matenda<sup>1\*</sup>, Nkongolo Bernard-Kennedy<sup>2</sup>,

Marc Bosonkie<sup>2</sup> and Nelly Dikamba<sup>3</sup>

<sup>1</sup>MD, MPH – Department of Community Health, School of Public Health, University of Kinshasa (UNIKIN), Kinshasa, Democratic Republic of the Congo.

<sup>2</sup>MD, MPH – Department of Nutrition, School of Public Health, University of Kinshasa, (UNIKIN), Kinshasa, Democratic Republic of the Congo.

<sup>3</sup>MD, MPH, PhD – Department of Biostatistics, School of Public Health, University of Kinshasa (UNIKIN), Kinshasa, Democratic Republic of the Congo.

### Article Info:

#### \*Corresponding author:

**Théophile Matenda,**

MD, MPH – Department of Community Health, School of Public Health, University of Kinshasa (UNIKIN), Kinshasa, Democratic Republic of the Congo.

[theophilematenda5@gmail.com](mailto:theophilematenda5@gmail.com)

DOI: [10.68819/anslhs.2026.0003](https://doi.org/10.68819/anslhs.2026.0003)

[Received 27-08-2026,

Accepted 11-10-2026,

Published 23-10-2026]

**Publishers Note:** Natural Science Association stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Copyright:** ©2026 by the author(s). This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 license (unless stated otherwise) permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

<https://creativecommons.org/licenses/by/4.0/>

### Abstract

**Introduction:** Measles remains a major cause of childhood morbidity and mortality in Africa. In the DRC, recurrent outbreaks reflect insufficient and uneven vaccine coverage, especially in rural areas. This study aimed to identify factors associated with measles resurgence among children aged 6 months to 10 years in Ludimbi Lukula Health Zone from 2016 to 2022.

**Methods:** A case-control study was conducted with 300 measles cases and 600 controls. Data were collected via KoboCollect from parents. Variables included vaccination status, nutritional condition, medical history, living conditions, and education level. Chi-square tests and logistic regression were used for analysis.

**Results:** Cases were concentrated in the Bemane (23.7%) and Luanga (19.7%) areas. Acute malnutrition was more prevalent among children with measles (54.3%) compared to those without (49.0%), with increased severity in boys and infants under one year old. Children from poor households were particularly vulnerable (35.0%). More than 90% of mothers were aware of measles, primarily through health workers, and considered vaccination the main form of prevention. Several factors were identified: health area, education, occupation, financial resources, age, and vaccination status, with the risk being four times higher among unvaccinated children.

**Conclusion:** Measles occurrence was associated with nutritional, socioeconomic, vaccination, and health-system-related factors. Integrated public health actions are needed to strengthen vaccine coverage, improve nutritional care, and include vulnerable populations in prevention strategies.

**Keywords:** Measles, associated factors, children, vaccination, Ludimbi Lukula

**Cite this article:** Théophile Matenda, Nkongolo Bernard-Kennedy, Marc Bosonkie and Nelly Dikamba, 2026. Factors Associated with Measles Occurrence among Children Aged 6 Months to 10 Years in Ludimbi Health Zone, Lukula, Democratic Republic of the Congo, 2016–2022: A Case-Control Study. Advanced Natural Sciences: Life and Health Sciences, 3(2), 194-209 <https://doi.org/10.68819/anslhs.2026.0003>

## I. Introduction

Measles is a highly contagious acute viral infection, primarily affecting children, with no specific cure. It manifests as a feverish rash accompanied by respiratory symptoms [1]. The introduction of an effective and inexpensive vaccine in the 1960s led to a significant reduction in global morbidity. However, the disease persists in countries with low vaccination coverage, particularly in Africa and Asia, where it causes large-scale outbreaks [1]. In 2019, the WHO reported a 300% increase in global measles cases in the first quarter compared to 2018, with a 700% increase in Africa. By that date, 170 countries had reported 112,163 cases, compared to 28,124 cases in 163 countries the previous year [2]. In Africa, nearly 17,500 cases were recorded between January and March 2022, an increase of 400% compared to the same period in 2021. Twenty countries reported outbreaks in the first quarter of 2022, compared to twelve in 2021 [3,4].

The Democratic Republic of Congo (DRC) experiences recurring measles outbreaks every two to three years, with peaks of 182,485 cases in 2005, 133,802 in 2011, and 88,381 in 2013. In 2018, 100 health zones were affected, with 124 more zones added in the first half of 2019, followed by 129 in the second half. In total, 353 out of 519 zones experienced a confirmed outbreak over two years [5]. Between 2019 and 2020, the DRC suffered the world's largest measles epidemic, with more than 380,000 cases and 7,000 deaths, primarily among children under five [6,7]. In 2023, Lomami province reported 20,791 cases, including 600 infant deaths, and the Ludimbi health zone Lukula recorded 1,142 cases [8,9]. Despite vaccination campaigns, 21 of the 26 provinces reported cases in 2023, revealing the persistence of the problem [7]. Obstacles include geographical inaccessibility, weak health infrastructure, malnutrition, and a lack of awareness about the importance of vaccination [10].

Ludimbi health zone In Lukula , several factors are implicated in the worsening situation: poor cold chain coverage, lack of training for

vaccination providers, insufficient parental awareness, poverty, overcrowding, and malnutrition [11, 12, 13, 14]. The strategies implemented—improving cold chain equipment, providing EPI training, and increasing the number of outreach and mobile vaccination sessions—have had a limited impact [13]. Despite a reduction in cases in some areas, measles persists in Ludimbi. Lukula , due to unfavorable socio-economic conditions and inadequate dietary practices [10]. To effectively combat this preventable disease, it is essential to strengthen vaccination campaigns, improve access to care, and intensify community awareness. An integrated approach combining routine immunization, nutrition, and health infrastructure is indispensable. This study aims to identify the factors associated with the resurgence of measles cases in children aged 6 months to 10 years in the Ludimbi health zone. Lukula , in order to guide future interventions.

## II. Methodology

### Design

We conducted a case-control study with children aged 6 months to 10 years living in the Ludimbi health zone Lukula .

### Study framework

Ludimbi health zone Lukula , located 100 km from Kabinda in Lomami province , covers 11,800 km<sup>2</sup> and has a low population density of nearly 200,000 inhabitants. It has a General Referral Hospital, 15 health centers, and 11 functional storage facilities. Despite a structured healthcare system, access to care remains difficult in remote areas such as Kitengie, Kiasame , and Kifuenkese . The population relies primarily on subsistence farming and small-scale livestock breeding, in a context of economic insecurity and limited access to essential services. These conditions exacerbate health vulnerabilities, particularly among children.

### Sampling

#### a) Population

The sampling for this case-control study focuses on children aged 6 months to 10 years residing

in the Ludimbi health zone Lukula, Lomami Province. Participants were selected from health areas that had recorded cases of measles during the study period, and mothers or caregivers served as respondents to provide the necessary information. Cases included children diagnosed with measles, whether they tested positive or not, and who had attended a local health facility. Controls were children of the same age who had not contracted measles, living in the same villages as the cases and sharing similar environmental conditions. Excluded were children with measles who were unavailable at the time of the survey, as well as controls whose parents or caregivers refused to provide informed consent.

#### b) Size

The study focused on the six health areas that recorded the highest number of measles cases out of the fifteen in the Ludimbi health zone. Lukula. The calculation of the sample size was carried out using the StatCalc program of the Epi Info 7 software, based on the parameters of a study carried out in Mali by Diarra Moussa Modibo et al.[1], with a confidence level of 95%, a power of 80%, a proportion of exposed controls of 50% and an odds ratio of 2.55.

$$n \geq 2 \times (Z\alpha + Z1 - \beta)^2 \times p(1-p) / (p_0 - p_1)^2 n$$

After inserting the values into the formula:

$$n \geq 2 \times (1.96 + 0.84)^2 \times 0.085 \times (1 - 0.085) / (0.05 - 0.12)^2 n$$

This gives:  $n \geq 272.5 \approx 273$  cases.

This calculation allowed us to estimate the number of cases needed at 273, corrected by a factor of 0.75 to obtain 410 controls. Thus, the calculated sample size was 615, comprising 273 cases and 410 controls. Due to organizational factors in the six selected health areas and to increase the study's statistical power, the number of cases was adjusted to 300 and the number of controls to 600, bringing the final sample size to 900.

#### c) Sampling technique

Cases were selected systematically from the line lists of children diagnosed with measles in the

six most affected health areas. Each eligible case was assigned a sequential number from 1 to N. The sampling interval (k) was calculated by dividing the total number of eligible cases by the required sample size. A random starting point was selected between 1 and k, after which subsequent cases were selected at regular intervals. In the event of refusal to participate or failure to meet the eligibility criteria, a replacement was selected using the same systematic sampling procedure, where feasible.

Controls were selected using a non-probability sampling approach based on geographical proximity to the identified cases. Eligible children without a history of measles were approached in nearby households, starting with plots adjacent to those of the cases. If no eligible control was available in the immediate vicinity, the search was extended to other nearby plots within the same health area. No individual or frequency matching was performed on the basis of age or sex. This approach was adopted for operational feasibility; however, it does not guarantee that the controls are representative of the population from which the cases arose. The potential for selection bias was therefore acknowledged when interpreting the findings.

#### Study variables

The variables studied in this research include a dependent variable—measles—defined as the presence of febrile maculopapular rash with or without rhinorrhea and conjunctivitis, or confirmed by laboratory testing, coded dichotomously (1: yes, 0: no). Independent variables include socioeconomic factors (income level and daily expenditure), demographic factors (child's sex and age, mother's marital status, head of household's occupation), educational factors (respondent's level of education), environmental factors (living environment, overcrowding), nutritional factors (nutritional status based on mid-upper arm circumference), and health factors (measles vaccination status). These variables were measured using nominal or ordinal scales, allowing for the exploration of their association with the occurrence of measles in children aged

6 months to 10 years in the Ludimbi health zone. Lukula .

### **Data collection**

Data collection was carried out using structured interviews administered to the parents of children with measles and to the parents of control subjects, selected according to predefined criteria in neighboring villages. A detailed questionnaire, designed and deployed via the Kobocollect mobile platform, enabled secure, real-time digital data entry, reducing errors and facilitating data processing. Before the start of the study, a pre-test was conducted to validate the tools, and the interviewers, healthcare professionals fluent in the local language, were trained on the study objectives, interview techniques, ethics, confidentiality, and the use of Kobocollect. Field supervision ensured quality control through regular monitoring, immediate corrections, and debriefing meetings. Following approval from the ethics committee, contacts were established with academic and health authorities and local training managers to obtain the necessary authorizations and lists. Field visits were planned to cover the targeted health areas, with logistical support including transportation, equipment, and data management. This rigorous and structured process ensured the reliability and ethical compliance of the data collection.

### **Data Processing and Analysis**

After data collection, the data were exported from Kobocollect and analyzed using SPSS version 25. Categorical variables were analyzed using absolute and relative frequencies, while quantitative variables were summarized using measures of central tendency and dispersion, according to their Kolmogorov-Smirnov distribution. In this study, nutritional status was assessed using mid-upper arm circumference (MUAC) and age to calculate MUAC-for-age Z-scores using WHO Anthro software for children under 5 years and WHO AnthroPlus for children aged 5 years and older. The use of these two software packages followed WHO recommendations according to the child's age group. Nutritional status was classified

according to the applicable WHO reference standards. A Z-score below  $-2$  was considered indicative of a value below the reference range (global acute malnutrition (GAM)), whereas a Z-score of  $-2$  or higher indicated a value within the reference range. Associations between variables were explored using the Chi-square test, and binary logistic regression was used to identify factors related to the occurrence of measles, with calculation of odds ratios and 95% confidence intervals.

### **Ethical considerations**

The study protocol was reviewed and approved by the Ethics Committee of the Kinshasa School of Public Health (approval number: CE/ESP/178/2025) in 2025. The study involved a secondary analysis of data extracted from health facility records and case-identification forms concerning children registered in healthcare facilities between 2016 and 2022. Thus, the ethical approval obtained in 2025 related to the present analysis of previously collected data, whereas the data covered the period from 2016 to 2022. The confidentiality and anonymity of the information were maintained throughout data extraction, processing and analysis. Access to the records was restricted to the research team, and identifying information was handled in accordance with applicable ethical requirements. The study was conducted in accordance with the principles of respect for persons, confidentiality and responsible use of health data.

## **Results**

### **Socio-demographic and socio-economic characteristics of households**

Analysis of the table shows a varied geographical distribution of respondents according to health area. Cases are mainly concentrated in the Bemane (23.7%) and Luanga (19.7%) health areas. Sociodemographically, the majority of respondents are women, aged 15 to 49, and primarily mothers of the affected children. Primary and secondary education levels dominate, with 44.3% of cases and 42.7 % of controls having a secondary education. More than half of the households surveyed in

both groups have fewer than six members. Economically, 82.5% of controls live on less than 3,000 FC per day per person, compared to 66.3% of cases. Finally, 41% of cases have a low socioeconomic level, while 35.8% of controls have a middle socioeconomic level. (Table I)

#### **Nutritional status of children aged 6 months to 10 years in surveyed households**

The results in Table II show a higher proportion of acute malnutrition among children with measles (54.3%) compared to controls (49.0%). According to Table III, the analysis aimed to determine whether nutritional status was associated with age, sex, and household wealth level within each study group. No statistically significant association was observed with sex among cases ( $p = 0.87$ ) or controls ( $p = 0.058$ ), while age was significantly associated with nutritional status among controls ( $p = 0.006$ ). Household wealth showed no statistically significant association, although children from poor households appeared more vulnerable, with 52.8% affected.

#### **Knowledge and perceptions of women and childcare providers regarding vaccination and/or measles**

The data reveal that the majority of respondents, both among cases (96.3%) and controls (91.7%), had heard of measles, primarily from healthcare workers. The most recognized symptoms were rash, fever, and cough. Over 80% of participants knew of at least one means of protection, with vaccination widely cited as the primary method (73.3% among cases, 71.3% among controls) (Table IV). The majority of respondents, both among cases (82.7%) and controls (79.7%), consider the vaccination to be effective against measles. A very positive perception of the vaccination is observed, with over 97% rating it as "good" among cases and 98.5% among controls. Finally, over 94.7% of cases, compared to 96.3% of controls, believe that the vaccine effectively protects the child (Table V).

#### **Children's vaccination status**

The majority of children, both cases and controls, had been vaccinated against measles (96.7% and 98.7%, respectively). However, a

slightly higher proportion of affected children (3.3%) had not been vaccinated compared to controls (1.3%), which, although marginal, could contribute to the occurrence of cases in a context of incomplete vaccination coverage (Figure 1).

#### **Factors associated with the occurrence of measles cases in children aged 6 months to 10 years**

Bivariate analysis using the chi-square test showed several significant associations ( $p < 0.05$ ) with the occurrence of measles. Cases varied according to the health area ( $p = 0.001$ ) and were more frequent in children aged 6–11 months and  $\geq 60$  months ( $p = 0.001$ ). The respondent's secondary education level ( $p = 0.043$ ) and teaching occupation ( $p = 0.01$ ) were associated with a higher incidence. Households with an expenditure  $\geq 3000$  FC/day had more cases ( $p = 0.001$ ). Lack of information about measles ( $p = 0.009$ ) and lack of vaccination, particularly against measles ( $p = 0.013$ ), were also aggravating factors. (Table VI)

Bivariate and multivariate logistic regression analyses identified several factors significantly associated with measles occurrence. Children residing in the Bemane (adjusted odds ratio [aOR] = 9.42) and Kiasame (aOR = 4.96) health areas had higher odds of measles occurrence than those living in Bashimikie. Similarly, having a respondent with secondary education (aOR = 1.67) and having a teacher as the respondent (aOR = 2.01) were associated with higher odds of measles occurrence. Children from households with daily expenditures of  $\geq 3,000$  Congolese francs (FC) also had higher odds of measles occurrence (aOR = 2.18). Conversely, children whose respondents had never heard of measles had lower odds of measles occurrence (aOR = 0.45). Compared with children aged  $\geq 60$  months, those aged 12–23 months (aOR = 0.11) and 24–59 months (aOR = 0.63) had lower odds of measles occurrence. Finally, children who had not been vaccinated against measles had higher odds of measles occurrence (aOR = 4.23, 95% CI: 1.22–14.63), highlighting the importance of measles vaccination.

**Table I:** Distribution according to the sociodemographic and socioeconomic characteristics of respondents

| Variables                                                           | Case  |      | Control |      |
|---------------------------------------------------------------------|-------|------|---------|------|
|                                                                     | n=300 | %    | n=600   | %    |
| <b>Health area</b>                                                  |       |      |         |      |
| <b>Bashimikie</b>                                                   | 51    | 17.0 | 102     | 17.0 |
| <b>Bemane</b>                                                       | 71    | 23.7 | 142     | 23.7 |
| <b>Kíasame</b>                                                      | 49    | 16.3 | 98      | 16.3 |
| <b>Luanga</b>                                                       | 59    | 19.7 | 118     | 19.7 |
| <b>Miombe 2</b>                                                     | 12    | 4.0  | 24      | 4.0  |
| <b>Miombe muavi</b>                                                 | 58    | 19.3 | 116     | 19.3 |
| <b>Respondent's gender</b>                                          |       |      |         |      |
| <b>Female</b>                                                       | 188   | 62.7 | 406     | 67.7 |
| <b>Male</b>                                                         | 112   | 37.3 | 194     | 32.3 |
| <b>Respondent's age (in years)</b>                                  |       |      |         |      |
| <b>&lt; 18</b>                                                      | 4     | 1.3  | 8       | 1.3  |
| <b>18-35</b>                                                        | 115   | 38.3 | 266     | 44.3 |
| <b>36-50</b>                                                        | 128   | 42.7 | 233     | 38.8 |
| <b>&gt; 50</b>                                                      | 53    | 17.7 | 93      | 15.5 |
| <b>Relationship with the child</b>                                  |       |      |         |      |
| <b>Other relatives (brother, sister, aunt, uncle, grandparents)</b> | 16    | 5.3  | 44      | 7.3  |
| <b>Mother</b>                                                       | 184   | 61.3 | 389     | 64.8 |
| <b>Father</b>                                                       | 93    | 31.0 | 157     | 26.2 |
| <b>Tutor</b>                                                        | 7     | 2.3  | 10      | 1.7  |
| <b>Respondent's level of education</b>                              |       |      |         |      |
| <b>No education</b>                                                 | 34    | 11.3 | 100     | 16.7 |
| <b>Primary</b>                                                      | 114   | 38.0 | 256     | 42.7 |
| <b>Secondary</b>                                                    | 149   | 49.6 | 237     | 39.5 |
| <b>Superior</b>                                                     | 3     | 1.0  | 7       | 1.2  |
| <b>Number of people in the household</b>                            |       |      |         |      |
| <b>&lt;6</b>                                                        | 167   | 55.7 | 308     | 51.3 |
| <b>&gt;=6</b>                                                       | 133   | 44.3 | 292     | 48.7 |
| <b>Daily household expenses/day</b>                                 |       |      |         |      |
| <b>Less than 3000 FC/Day/person</b>                                 | 199   | 66.3 | 495     | 82.5 |
| <b>Greater than or equal to 3000 FC/Day/person</b>                  | 101   | 33.7 | 105     | 17.5 |
| <b>Residence status in the plot</b>                                 |       |      |         |      |
| <b>Other (housed, family, etc. )</b>                                | 2     | 0.7  | 2       | 0.3  |
| <b>Tenant</b>                                                       | 53    | 17.7 | 82      | 13.7 |
| <b>Owner</b>                                                        | 245   | 81.7 | 516     | 86.0 |
| <b>Wealth index</b>                                                 |       |      |         |      |
| <b>Low</b>                                                          | 123   | 41.0 | 211     | 35.2 |
| <b>Middle</b>                                                       | 85    | 28.3 | 215     | 35.8 |
| <b>High</b>                                                         | 92    | 30.7 | 174     | 29.0 |

**Table II.** Nutritional status of the child measured by mid-upper arm circumference according to age (Z-score for age)

| Nutritional status           | Case (n = 300) | Controls (n = 600) |
|------------------------------|----------------|--------------------|
| Good (No acute malnutrition) | 137 (45.7)     | 306 (51.0)         |
| Bad / MAG                    | 163 (54.3)     | 294 (49.0)         |
| Total                        | 300 (100,0)    | 600 (100,0)        |

$p(X^2)=0,131$

**Table III:** Distribution of children's nutritional status according to sex, age and level of wealth ownership

| Features                | Case               |               |      | Control            |               |       |
|-------------------------|--------------------|---------------|------|--------------------|---------------|-------|
|                         | Nutritional status |               |      | Nutritional status |               |       |
|                         | Good n (%)         | Bad/MAG n (%) | p    | Good n (%)         | Bad/MAG n (%) | p     |
| <b>Sex of the child</b> |                    |               |      |                    |               |       |
| <b>Female</b>           | 66 (45.2)          | 80 (54.8)     | 0.87 | 141 (55.5)         | 113 (44.5)    | 0.058 |
| <b>Male</b>             | 71 (46.1)          | 83 (53.9)     |      | 165 (47.7)         | 181 (52.3)    |       |
| <b>Child's age</b>      |                    |               |      |                    |               |       |
| <b>6–11 months</b>      | 6 (60.0)           | 4 (40.0)      | 0.81 | 10 (83.3)          | 2 (16.7)      | 0.006 |
| <b>12–23 months</b>     | 1 (50.0)           | 1 (50.0)      |      | 26 (72.2)          | 10 (27.8)     |       |
| <b>24–59 months</b>     | 47 (46.1)          | 55 (53.9)     |      | 127 (49.0)         | 133 (51.0)    |       |
| <b>≥60 months</b>       | 83 (44.6)          | 103 (55.4)    |      | 143 (49.0)         | 149 (51.0)    |       |
| <b>Wealth index</b>     |                    |               |      |                    |               |       |
| <b>Low</b>              | 58 (47.2)          | 65 (52.8)     | 0.86 | 118 (55.9)         | 93 (44.1)     | 0.16  |
| <b>Middle</b>           | 39 (45.9)          | 46 (54.1)     |      | 107 (49.8)         | 108 (50.2)    |       |
| <b>High</b>             | 40 (43.5)          | 52 (56.5)     |      | 81 (46.6)          | 93 (53.4)     |       |

**Table IV:** Mothers'/Carers' Knowledge of Measles

| Respondents' knowledge and perceptions about measles | Case  |      | control |      |
|------------------------------------------------------|-------|------|---------|------|
|                                                      | n=300 | %    | n=600   | %    |
| <b>Have you ever heard of measles?</b>               |       |      |         |      |
| <b>No</b>                                            | 11    | 3.7  | 50      | 8.3  |
| <b>Yes</b>                                           | 289   | 96.3 | 550     | 91.7 |
| <b>Source of information</b>                         |       |      |         |      |
| <b>Posters</b>                                       | 4     | 1.3  | 38      | 6.3  |
| <b>Other (Social networks, internet):</b>            | 10    | 3.3  | 17      | 2.8  |
| <b>Healthcare personnel</b>                          | 205   | 68.3 | 366     | 61.0 |
| <b>Radio</b>                                         | 65    | 21.7 | 118     | 19.7 |
| <b>Neighbors/friends</b>                             | 16    | 5.3  | 61      | 10.2 |
| <b>Known signs of measles</b>                        |       |      |         |      |
| <b>Fever</b>                                         | 206   | 68.7 | 403     | 67.2 |
| <b>Cough</b>                                         | 112   | 37.3 | 162     | 27   |
| <b>Rash</b>                                          | 242   | 80.7 | 503     | 83.8 |
| <b>Runny nose</b>                                    | 92    | 30.7 | 117     | 19.5 |
| <b>Bright red spots</b>                              | 91    | 30.3 | 115     | 19.2 |

| Knowledge of means of protection (1/2)                      |     |      |     |      |
|-------------------------------------------------------------|-----|------|-----|------|
| No                                                          | 53  | 17.7 | 114 | 19.0 |
| Yes                                                         | 247 | 82.3 | 486 | 81.0 |
| Knowledge of means of protection (2/2)                      |     |      |     |      |
| Other ( Isolation of cases, health education, surveillance) | 1   | 0.3  | 4   | 0.7  |
| Good nutrition                                              | 9   | 3.0  | 27  | 4.5  |
| Timely medical care                                         | 17  | 5.7  | 26  | 4.3  |
| Vaccination                                                 | 220 | 73.3 | 428 | 71.3 |

Table V. Mothers'/Carers' Perceptions of Measles

| Respondents' Perceptions                               | Case  |      | Control |      |
|--------------------------------------------------------|-------|------|---------|------|
|                                                        | n=300 | %    | n=600   | %    |
| Do you think vaccination is effective against measles? |       |      |         |      |
| Don't know                                             | 47    | 15.7 | 106     | 17.7 |
| No                                                     | 5     | 1.7  | 16      | 2.7  |
| Yes                                                    | 248   | 82.7 | 478     | 79.7 |
| What are your thoughts on vaccination?                 |       |      |         |      |
| Other (don't know, no choice)                          | 4     | 1.3  | 4       | 0.7  |
| Good                                                   | 291   | 97   | 591     | 98.5 |
| Bad .                                                  | 5     | 1.7  | 5       | 0.8  |
| Do you think the vaccine can protect the child?        |       |      |         |      |
| No                                                     | 16    | 5.3  | 22      | 3.7  |
| Yes                                                    | 284   | 94.7 | 578     | 96.3 |




Figure 1: Child's vaccination status

**Table VI:** Factors associated with the occurrence of measles cases in children aged 6 months to 10 years in the Ludimbi Health Zone Lukula according to the Chi-square test

| Variables                                  | Case        | Control     | Total       | p(X <sup>2</sup> ) |
|--------------------------------------------|-------------|-------------|-------------|--------------------|
| <b>Health area</b>                         |             |             |             | 0.001              |
| <b>Bashimikie</b>                          | 51 (17.0%)  | 102 (17.0%) | 153 (17.0%) |                    |
| <b>Bemane</b>                              | 71 (23.7%)  | 142 (23.7%) | 213 (23.7%) |                    |
| <b>Kîasame</b>                             | 49 (16.3%)  | 98 (16.3%)  | 147 (16.3%) |                    |
| <b>Luanga</b>                              | 59 (19.7%)  | 118 (19.7%) | 177 (19.7%) |                    |
| <b>Miombe Etala</b>                        | 12 (4.0%)   | 24 (4.0%)   | 36 (4.0%)   |                    |
| <b>Miombe Muavi</b>                        | 58 (19.3%)  | 116 (19.3%) | 174 (19.3%) |                    |
| <b>Respondent's age (years)</b>            |             |             |             | 0.072              |
| <b>&lt; 18</b>                             | 4 (33.3%)   | 8 (66.7%)   | 12          |                    |
| <b>18–35</b>                               | 115 (30.2%) | 266 (69.8%) | 381         |                    |
| <b>36–50</b>                               | 128 (35.5%) | 233 (64.5%) | 361         |                    |
| <b>&gt; 50</b>                             | 53 (36.3%)  | 93 (63.7%)  | 146         |                    |
| <b>Relationship with the child</b>         |             |             |             | 0.29               |
| <b>Mother</b>                              | 184 (32.1%) | 389 (67.9%) | 573 (100%)  |                    |
| <b>Father</b>                              | 93 (37.2%)  | 157 (62.8%) | 250 (100%)  |                    |
| <b>Tutor</b>                               | 7 (41.2%)   | 10 (58.8%)  | 17 (100%)   |                    |
| <b>Other</b>                               | 16 (26.7%)  | 44 (73.3%)  | 60 (100%)   |                    |
| <b>Respondent's level of education</b>     |             |             |             | 0.043              |
| <b>Primary</b>                             | 114 (30.8%) | 256 (69.2%) | 370 (100%)  |                    |
| <b>Secondary</b>                           | 149 (38.6%) | 237 (61.4%) | 386 (100%)  |                    |
| <b>Superior</b>                            | 3 (30.0%)   | 7 (70.0%)   | 10 (100%)   |                    |
| <b>Other</b>                               | 16 (37.2%)  | 27 (62.8%)  | 43 (100%)   |                    |
| <b>Respondent's profession</b>             |             |             |             | 0.01               |
| <b>Farmer/Livestock Breeder</b>            | 230 (31.8%) | 493 (68.2%) | 723 (100%)  |                    |
| <b>Teacher</b>                             | 38 (51.4%)  | 36 (48.6%)  | 74 (100%)   |                    |
| <b>Shopkeeper</b>                          | 3 (25.0%)   | 9 (75.0%)   | 12 (100%)   |                    |
| <b>Other</b>                               | 29 (31.9%)  | 62 (68.1%)  | 91 (100%)   |                    |
| <b>Marital status of mother/guardian</b>   |             |             |             | 0.6                |
| <b>Bride</b>                               | 287 (33.8%) | 563 (66.2%) | 850 (100%)  |                    |
| <b>Bachelor</b>                            | 3 (27.3%)   | 8 (72.7%)   | 11 (100%)   |                    |
| <b>Divorcee</b>                            | 8 (29.6%)   | 19 (70.4%)  | 27 (100%)   |                    |
| <b>Widow</b>                               | 2 (18.2%)   | 9 (81.8%)   | 11 (100%)   |                    |
| <b>Free union</b>                          | 0 (0.0%)    | 1 (100.0%)  | 1 (100%)    |                    |
| <b>Number of people in the household</b>   |             |             |             | 0.22               |
| <b>&lt;6 people</b>                        | 167 (35.2%) | 308 (64.8%) | 475 (100%)  |                    |
| <b>6 people or more</b>                    | 133 (31.3%) | 292 (68.7%) | 425 (100%)  |                    |
| <b>Daily household expenditure per day</b> |             |             |             | 0.001              |
| <b>&lt;3000 FC</b>                         | 199 (28.7%) | 495 (71.3%) | 694 (100%)  |                    |
| <b>≥3000 FC</b>                            | 101 (49.0%) | 105 (51.0%) | 206 (100%)  |                    |
| <b>Wealth index</b>                        |             |             |             | 0.06               |
| <b>Low</b>                                 | 123 (36.8%) | 211 (63.2%) | 334 (100%)  |                    |
| <b>Middle</b>                              | 85 (28.3%)  | 215 (71.7%) | 300 (100%)  |                    |
| <b>High</b>                                | 92 (34.6%)  | 174 (65.4%) | 266 (100%)  |                    |
| <b>Have you ever heard of measles?</b>     |             |             |             | 0.009              |
| <b>Yes</b>                                 | 289 (34.4%) | 550 (65.6%) | 839 (100%)  |                    |

|                                                             |             |             |            |       |
|-------------------------------------------------------------|-------------|-------------|------------|-------|
| No                                                          | 11 (18.0%)  | 50 (82.0%)  | 61 (100%)  |       |
| <b>Sex of the child</b>                                     |             |             |            | 0.07  |
| Female                                                      | 146 (36.5%) | 254 (63.5%) | 400 (100%) |       |
| Male                                                        | 154 (30.8%) | 346 (69.2%) | 500 (100%) |       |
| <b>Child's age (in months)</b>                              |             |             |            | 0.001 |
| 6–11                                                        | 10 (45.5%)  | 12 (54.5%)  | 22 (100%)  |       |
| 12–23                                                       | 2 (5.3%)    | 36 (94.7%)  | 38 (100%)  |       |
| 24–59                                                       | 102 (28.2%) | 260 (71.8%) | 362 (100%) |       |
| ≥60                                                         | 186 (38.9%) | 292 (61.1%) | 478 (100%) |       |
| <b>Child's nutritional status ( Z-score) Muac for age )</b> |             |             |            | 0.13  |
| Good (No acute malnutrition)                                | 137 (30.9)  | 306 (69.1)  | 443 (100)  |       |
| Bad (MAG)                                                   | 163 (35.7)  | 294 (64.3)  | 457 (100)  |       |
| <b>The child has already been vaccinated.</b>               |             |             |            | 0.043 |
| Yes                                                         | 290 (32.9%) | 592 (67.1%) | 882 (100%) |       |
| No                                                          | 10 (55.6%)  | 8 (44.4%)   | 18 (100%)  |       |
| <b>The child was vaccinated against measles.</b>            |             |             |            |       |
| Yes                                                         | 291 (32.8%) | 596 (67.2%) | 887 (100%) | 0.013 |
| No                                                          | 9 (69.2%)   | 4 (30.8%)   | 13 (100%)  |       |

**Table VII:** Factors associated with the occurrence of measles cases in children aged 6 months to 10 years in the Ludimbi Health Zone Lukula in bivariate and multivariate logistic regression

| Variables                              | Case<br>n=300 | Control<br>n=600 | ORb<br>(95% CI)    | P      | ORa<br>(95% CI)   | P      |
|----------------------------------------|---------------|------------------|--------------------|--------|-------------------|--------|
| <b>Health area</b>                     |               |                  |                    |        |                   |        |
| Bashimikie                             | 51 (34.0%)    | 99 (66.0%)       | 1                  |        | 1                 |        |
| Bemane                                 | 71 (48.0%)    | 77 (52.0%)       | 10.91 (5.65–21.07) | <0.001 | 9.42 (4.89–18.14) | <0.001 |
| Kiasame                                | 49 (32.9%)    | 100 (67.1%)      | 5.80 (2.94–11.45)  | <0.001 | 4.96 (2.42–10.17) | <0.001 |
| Luanga                                 | 59 (39.9%)    | 89 (60.1%)       | 7.84 (4.02–15.29)  | <0.001 | 1.42 (0.63–3.17)  | 0.392  |
| Miombe Etala                           | 12 (7.8%)     | 142 (92.2%)      | 6.14 (3.11–12.12)  | <0.001 | 1.10 (0.54–2.23)  | 0.45   |
| Miombe Muavi                           | 58 (38.4%)    | 93 (61.6%)       | 7.38 (3.78–14.41)  | <0.001 | 1.41 (0.74–2.68)  | 0.8    |
| <b>Respondent's level of education</b> |               |                  |                    |        |                   |        |
| None                                   | 34 (25.4%)    | 100 (74.6%)      | 1                  |        | 1                 |        |
| Primary                                | 114 (30.8%)   | 256 (69.2%)      | 1.30 (0.82–2.07)   | 0.26   | 1.21 (0.74–1.98)  | 0.45   |
| Secondary                              | 149 (38.6%)   | 237 (61.4%)      | 1.80 (1.13–2.87)   | 0.014  | 1.67 (1.03–2.71)  | 0.038  |
| Superior                               | 3 (30.0%)     | 7 (70.0%)        | 1.26 (0.30–5.35)   | 0.75   | 0.74(0.39-1.45)   | 0.39   |
| <b>Respondent's profession</b>         |               |                  |                    |        |                   |        |
| Farmer/Livestock Breeder               | 230 (31.8%)   | 493 (68.2%)      | 1                  |        | 1                 |        |
| Teacher                                | 38 (51.4%)    | 36 (48.6%)       | 2.23 (1.39–3.58)   | 0.001  | 2.01 (1.21–3.36)  | 0.007  |

|                                                  |             |             |                   |        |                   |        |
|--------------------------------------------------|-------------|-------------|-------------------|--------|-------------------|--------|
| <b>Shopkeeper</b>                                | 3 (25.0%)   | 9 (75.0%)   | 0.71 (0.19–2.65)  | 0.61   | 2.02(0.85 – 4.79) | 0.107  |
| <b>Other</b>                                     | 29 (31.9%)  | 62 (68.1%)  | 1.01 (0.64–1.59)  | 0.96   | 1.24(0.69 – 2.22) | 0.46   |
| <b>Daily household expenditure per day</b>       |             |             |                   |        |                   |        |
| <b>&lt;3000 FC</b>                               | 199 (28.7%) | 495 (71.3%) | 1                 |        | 1                 |        |
| <b>≥3000 FC</b>                                  | 101 (49.0%) | 105 (51.0%) | 2.39 (1.73–3.29)  | <0.001 | 2.18 (1.56–3.05)  | <0.001 |
| <b>Wealth index</b>                              |             |             |                   |        |                   |        |
| <b>Low</b>                                       | 123 (36.8%) | 211 (63.2%) | 1                 |        | 1                 |        |
| <b>Middle</b>                                    | 85 (28.3%)  | 215 (71.7%) | 0.68 (0.48–0.97)  | 0.033  | 0.74 (0.51–1.07)  | 0.11   |
| <b>High</b>                                      | 92 (34.6%)  | 174 (65.4%) | 0.89 (0.63–1.27)  | 0.52   | 0.94 (0.64–1.38)  | 0.76   |
| <b>Have you ever heard of measles?</b>           |             |             |                   |        |                   |        |
| <b>Yes</b>                                       | 289 (34.4%) | 550 (65.6%) | 1                 |        | 1                 |        |
| <b>No</b>                                        | 11 (18.0%)  | 50 (82.0%)  | 0.42 (0.21–0.84)  | 0.014  | 0.45 (0.22–0.93)  | 0.032  |
| <b>Sex of the child</b>                          |             |             |                   |        |                   |        |
| <b>Female</b>                                    | 146 (36.5%) | 254 (63.5%) | 1                 |        | 1                 |        |
| <b>Male</b>                                      | 154 (30.8%) | 346 (69.2%) | 0.76 (0.56–1.03)  | 0.07   | 0.81 (0.59–1.12)  | 0.2    |
| <b>Child's age (in months)</b>                   |             |             |                   |        |                   |        |
| <b>6–11</b>                                      | 10 (46.7%)  | 12 (53.3%)  | 1.18 (0.52–2.66)  | 0.69   | 1.16 (0.49–2.75)  | 0.73   |
| <b>12–23</b>                                     | 2 (5.3%)    | 36 (94.7%)  | 0.10 (0.02–0.43)  | 0.002  | 0.11 (0.03–0.47)  | 0.003  |
| <b>24–59</b>                                     | 102 (28.2%) | 260 (71.8%) | 0.61 (0.45–0.84)  | 0.002  | 0.63 (0.46–0.88)  | 0.006  |
| <b>≥60</b>                                       | 186 (38.9%) | 292 (61.1%) | 1                 |        | 1                 |        |
| <b>The child has already been vaccinated.</b>    |             |             |                   |        |                   |        |
| <b>Yes</b>                                       | 290 (32.9%) | 592 (67.1%) | 1                 |        | 1                 |        |
| <b>No</b>                                        | 10 (55.6%)  | 8 (44.4%)   | 2.55 (0.98–6.63)  | 0.053  | 2.31 (0.86–6.23)  | 0.097  |
| <b>The child was vaccinated against measles.</b> |             |             |                   |        |                   |        |
| <b>Yes</b>                                       | 291 (32.8%) | 596 (67.2%) | 1                 |        | 1                 |        |
| <b>No</b>                                        | 9 (69.2%)   | 4 (30.8%)   | 4.61 (1.39–15.27) | 0.013  | 4.23 (1.22–14.63) | 0.023  |

#### IV. Discussion

##### Key Results

Ludimbi Health Zone Lukula reveals a high concentration of measles cases among children aged 6 months to 10 years, particularly in the Bemane and Kiasame health areas . Analysis of mid-upper arm circumference shows a high

prevalence of acute malnutrition among affected children (54.3%) compared to 49.0% among unaffected children, with increased severity in boys and infants under one year old. Children from households with low levels of wealth are also more vulnerable (35.0%). The majority of mothers or caregivers (over 90%) are familiar

with measles, primarily through health workers, and recognize its main clinical signs. Vaccination is perceived as the essential means of prevention, although some express doubts about its effectiveness. Vaccination coverage remains high in both groups (96.7% in cases and 98.7% in controls). Statistical analysis identifies several factors associated with the occurrence of measles: health area, respondent's education level and occupation, daily household expenses, child's age, and vaccination status. Unvaccinated children have a fourfold increased risk.

### **Knowledge of mothers and caregivers regarding measles and vaccination**

In the Ludimbi Health Zone, Lukula, more than 90% of mothers and caregivers had heard of measles, primarily through healthcare workers. Respondents generally recognized the main clinical signs, including fever, cough and rash, and identified vaccination as the primary means of prevention. The importance of caregiver awareness, maternal education and access to vaccination services has also been documented in sub-Saharan Africa [15,16]. However, some respondents in Ludimbi expressed doubts about vaccine effectiveness, potentially reflecting personal experiences or concerns about breakthrough infections.

The prominent role of healthcare workers in providing information represents an opportunity to strengthen health education and address misconceptions. Nevertheless, adequate knowledge does not necessarily translate into complete vaccination. Practical barriers, including geographical distance, vaccine availability, indirect costs and missed opportunities during healthcare visits, may contribute to incomplete immunization. Evidence from the Democratic Republic of the Congo has highlighted disparities in measles vaccination coverage, supporting the need to combine health education with accessible and reliable vaccination services [17].

### **Factors associated with measles occurrence**

The findings from Ludimbi reflect challenges reported in other African settings, where geographical inequalities and socioeconomic circumstances may be associated with

differences in measles occurrence. Studies conducted in the DRC and systematic reviews from Africa have documented persistent challenges in vaccination coverage and measles control [18,19]. In the present study, children residing in Bemane (aOR = 9.42; 95% CI: 4.89–18.14) and Kiasame (aOR = 4.96; 95% CI: 2.42–10.17) had higher odds of measles occurrence than those living in Bashimikie. These substantial differences may reflect geographical inequalities in vaccination coverage, access to healthcare, vaccine availability, population mobility or outbreak detection. However, these mechanisms were not necessarily measured in the present study and should be considered possible explanations rather than established causes. Regional surveillance reports have documented persistent measles transmission and difficulties in achieving elimination targets in the WHO African Region [20]. Economic and occupational characteristics were also associated with measles occurrence in Ludimbi. Children from households reporting daily expenditure of at least 3,000 Congolese francs had higher odds of being cases (aOR = 2.18; 95% CI: 1.56–3.05). Similarly, having a respondent with secondary education (aOR = 1.67; 95% CI: 1.03–2.71) and having a teacher as the respondent (aOR = 2.01; 95% CI: 1.21–3.36) were associated with higher odds of case status. These findings should be interpreted cautiously because the observed associations may reflect differences in residence, mobility, household characteristics or other unmeasured factors. Socioeconomic inequalities in childhood vaccination have been documented in low- and middle-income countries, highlighting the importance of considering social and health-service contexts when interpreting these findings [21,22]. The present results do not establish that higher household expenditure, secondary education or teaching as an occupation directly increases measles susceptibility.

Nutritional vulnerability is another important consideration in measles prevention. Although nutritional status was not identified as a statistically significant factor in the present

analysis, this does not exclude its potential relevance to child health. Previous research has examined the epidemiology of measles outbreaks and the vulnerability of children in affected populations [18,19]. However, in Ludimbi, children aged 12–23 months (aOR = 0.11; 95% CI: 0.03–0.47) and 24–59 months (aOR = 0.63; 95% CI: 0.46–0.88) had lower odds of measles occurrence than children aged 60 months or older. This pattern differs from the commonly described vulnerability of younger children and warrants cautious interpretation, taking into account local vaccination patterns, exposure histories and the distribution of cases and controls across age groups.

From a behavioral perspective, concerns about vaccination and trust in health services may influence immunization uptake. The importance of supplementary immunization activities in reaching children missed by routine vaccination services has been documented in previous research [23]. In Ludimbi, respondents who had never heard of measles unexpectedly had lower odds of case status (aOR = 0.45; 95% CI: 0.22–0.93). This association should not be interpreted as a protective effect of limited awareness. The relatively small number of respondents in this category, potential selection or reporting bias, residual confounding, and possible measurement or coding problems may have influenced the result. Verification of the original data and variable coding is therefore advisable.

Finally, children who had not been vaccinated against measles had higher odds of measles occurrence (aOR = 4.23; 95% CI: 1.22–14.63). This association reinforces the importance of vaccination as a central component of measles prevention. National and regional immunization reports also emphasize the need to address vaccination gaps and strengthen strategies for reaching underserved populations [24]. Although the confidence interval indicates uncertainty regarding the magnitude of the association, the finding supports strengthening routine immunization, improving access to healthcare, addressing geographical disparities and implementing targeted community engagement. Because this was an observational

case-control study, these results identify associations rather than establish causal relationships.

### **Strengths and limitations of the study**

This study benefited from a case-control design involving 300 cases and 600 controls, systematic case selection, digital data collection using KoboCollect, and multivariable logistic regression. Nutritional status was assessed using MUAC-for-age Z-scores calculated with WHO Anthro for children under 5 years and AnthroPlus for those aged 5 years and older. Ethical approval was obtained from the relevant ethics committee. However, the final sample size differed from the initial calculation because of organizational considerations and the aim of improving statistical power. The non-probability selection of controls based on geographical proximity may have introduced selection bias and limits representativeness. The absence of documented vaccination histories may have caused recall bias, while social desirability may have influenced some responses. The study also did not adequately account for measles seasonality, and inaccessible areas may have limited generalizability. Finally, the observational design precludes causal conclusions, and the nutritional findings should be interpreted in relation to the specific anthropometric indicator used.

### **Conclusion**

This case-control study conducted in the Ludimbi Health Zone, Lukula, identified several factors associated with measles occurrence among children, including socioeconomic characteristics, vaccination status, and other health-related factors. Despite reported awareness of measles, gaps in vaccination and concerns about vaccine efficacy may remain. These findings highlight the importance of an integrated public health approach that strengthens vaccination coverage, promotes nutritional well-being, and improves access to healthcare services. Although causal relationships cannot be established from this observational design, the results provide useful

evidence to inform locally adapted measles prevention and control strategies.

**Acknowledgements:** We sincerely thank all the healthcare professionals who participated in this study, as well as the heads of the health facilities in the Ludimbi Health Zone. Lukula for their collaboration and availability.

**Disclaimer:** The opinions expressed in this manuscript are those of the authors and do not necessarily reflect those of associated health or academic institutions.

**Competing Interests:** The authors declare that they have no competing interests related to this study.

**Ethical statement:** This study was approved by the Ethics Committee of the Kinshasa School of Public Health (approval number: CE/ESP/178/2025). Informed consent was obtained from all participants before their inclusion in the study.

**Data availability:** The data used for this research are available from the corresponding author upon reasoned request. They are not publicly accessible in order to protect the confidentiality of the participants.

**Funding:** This research received no external funding. The costs associated with publishing this article were not covered by any third-party source.

**Author contributions:** Théophile Matenda: study design, supervision of data collection, analysis, manuscript writing.

- Nkongolo Bernard-Kennedy: contribution to the statistical analysis and critical review of the manuscript.
- Nelly Dikamba : Facilitation, methodological support, coordination and supervision of the writing, approval of the protocol and the manuscript

**Declaration of Generative AI and AI-Assisted Technologies:** Generative AI and AI-assisted tools were used to assist with language editing, grammar correction, and improvement of the clarity and readability of the manuscript. The authors reviewed and revised the AI-assisted content and take full responsibility for the accuracy, originality, integrity, and final content of the article. AI tools were not considered authors and were not assigned authorship.

## References

1. World Health Organization. Measles. Geneva: WHO; 2026 Jul 15 [cited 2026 Oct 11]. Available from: <https://www.who.int/news-room/factsheets/detail/measles>.
2. Moss WJ. Measles. Lancet. 2017;390(10111):2490–502. doi:10.1016/S0140-6736(17)31463-0.
3. World Health Organization. Measles vaccines: WHO position paper—April 2017. Wkly Epidemiol Rec. 2017;92(17):205–27.
4. Ashbaugh HR, Hoff NA, Doshi RH, Alfonso VH, Gadoth A, Mukadi P, et al. Predictors of measles vaccination coverage among children 6–59 months of age in the Democratic Republic of the Congo. Vaccine. 2018;36(4):587–93. doi:10.1016/j.vaccine.2017.11.049.
5. Chilot D, Belay DG, Shitu K, Gela YY, Getnet M, Mulat B, et al. Measles second dose vaccine utilization and associated factors among children aged 24–35 months in Sub-Saharan Africa: a multi-level analysis from recent DHS surveys. BMC Public Health. 2022;22:2070. doi:10.1186/s12889-022-14478-x.
6. Bobo FT, Asante A, Woldie M, Dawson A, Hayen A. Child vaccination in sub-Saharan Africa: increasing coverage addresses inequalities. Vaccine. 2022;40(1):141–50. doi:10.1016/j.vaccine.2021.11.005.
7. Tariku MK, Worede DT, Belete AH, Bante SA, Misikir SW. Attack rate, case fatality rate and determinants of measles infection during a measles outbreak in Ethiopia:

- systematic review and meta-analysis. *BMC Infect Dis.* 2023;23:756. doi:10.1186/s12879-023-08757-0.
8. Eshetu D, Tosisa W, Regassa BT, Hundie GB, et al. Epidemiology of measles outbreaks, incidence and associated risk factors in Ethiopia from 2000 to 2023: a systematic review and meta-analysis. *BMC Infect Dis.* 2024;24. doi:10.1186/s12879-024-09828-6.
  9. Masresha BG, Wiysonge CS, Katsande R, O'Connor PM, Lebo E, Perry RT. Tracking measles and rubella elimination progress—World Health Organization African Region, 2022–2023. *Vaccines (Basel).* 2024;12(8):949. doi:10.3390/vaccines12080949.
  10. World Health Organization. Guideline: vitamin A supplementation in infants and children 6–59 months of age. Geneva: WHO; 2011. Available from: <https://www.who.int/publications-detail-redirect/9789241501767>.
  11. World Health Organization. Vitamin A supplementation in infants and children 6–59 months of age [Internet]. Geneva: WHO; updated 2023 Aug 9. Available from: <https://www.who.int/tools/elena/interventions/vitamina-children>.
  12. Portnoy A, Jit M, HELLERINGER S, Verguet S. Impact of measles supplementary immunization activities on reaching children missed by routine programs. *Vaccine.* 2018;36(1):170–8. doi:10.1016/j.vaccine.2017.10.080.
  13. Portnoy A, Jit M, HELLERINGER S, Verguet S. Impact of measles supplementary immunization activities in low- and middle-income countries, 2000–2017. *BMC Med.* 2018;16:116. doi:10.1186/s12916-018-1120-6.
  14. Hajizadeh M. Socioeconomic inequalities in child vaccination in low/middle-income countries: what accounts for the differences? *J Epidemiol Community Health.* 2018;72(8):719–25. doi:10.1136/jech-2017-210296.
  15. Bobo FT, Asante A, Woldie M, Dawson A, Hayen A. Child vaccination in sub-Saharan Africa: increasing coverage addresses inequalities. *Vaccine.* 2022;40(1):141–150. doi:10.1016/j.vaccine.2021.11.005.
  16. Chilot D, Belay DG, Shitu K, Gela YY, Getnet M, Mulat B, et al. Measles second dose vaccine utilization and associated factors among children aged 24–35 months in Sub-Saharan Africa: a multi-level analysis from recent DHS surveys. *BMC Public Health.* 2022;22:2070. doi:10.1186/s12889-022-14478-x.
  17. Ashbaugh HR, Hoff NA, Doshi RH, Alfonso VH, Gadoth A, Mukadi P, et al. Predictors of measles vaccination coverage among children 6–59 months of age in Democratic Republic of the Congo. *Vaccine.* 2018;36(4):587–593. doi:10.1016/j.vaccine.2017.11.049.
  18. Tariku MK, Worede DT, Belete AH, Bante SA, Misikir SW. Attack rate, case fatality rate and determinants of measles infection during a measles outbreak in Ethiopia: systematic review and meta-analysis. *BMC Infect Dis.* 2023;23:756. doi:10.1186/s12879-023-08757-0.
  19. Eshetu D, Tosisa W, Regassa BT, Hundie GB, et al. Epidemiology of measles outbreaks, incidence and associated risk factors in Ethiopia from 2000 to 2023: a systematic review and meta-analysis. *BMC Infect Dis.* 2024. doi:10.1186/s12879-024-09828-6.
  20. Masresha BG, Wiysonge CS, Katsande R, O'Connor PM, Lebo E, Perry RT. Tracking measles and rubella elimination progress—WHO African Region, 2022–2023. *Vaccines (Basel).* 2024;12(8):949. doi:10.3390/vaccines12080949.
  21. Hajizadeh M. Socioeconomic inequalities in child vaccination in low/middle-income countries: what accounts for the differences? *J Epidemiol Community Health.* 2018;72(8):719–725. doi:10.1136/jech-2017-210296.

22. Bobo FT, Asante A, Woldie M, Dawson A, Hayen A. Child vaccination in sub-Saharan Africa: increasing coverage addresses inequalities. *Vaccine*. 2022;40(1):141–150. doi:10.1016/j.vaccine.2021.11.005.
23. Portnoy A, Jit M, Helleringer S, Verguet S. Impact of measles supplementary immunization activities on reaching children missed by routine programs. *Vaccine*. 2018;36(1):170–178. doi:10.1016/j.vaccine.2017.10.080.
24. World Health Organization Regional Office for Africa. Vaccine-preventable disease outbreaks on the rise in Africa. Brazzaville: WHO AFRO; 2022 Apr 28. Available from: <https://www.afro.who.int/news/vaccine-preventable-disease-outbreaks-rise-africa>.